

Model 1 Sebelum Dimoderasi Inflasi

Uji Normalitas

One-Sample Kolmogorov-Smirnov Test

		Unstandardized Residual	
N		75	
Normal Parameters ^{a,b}	Mean	.0000000	
	Std. Deviation	1.09385773	
Most Extreme Differences	Absolute	.184	
	Positive	.162	
	Negative	-.184	
Test Statistic		.184	
Asymp. Sig. (2-tailed) ^c		.000	
Monte Carlo Sig. (2-tailed) ^d	Sig.	.000	
	99% Confidence Interval	Lower Bound	.000
		Upper Bound	.000


a. Test distribution is Normal.

b. Calculated from data.

c. Lilliefors Significance Correction.

d. Lilliefors' method based on 10000 Monte Carlo samples with starting seed 1314643744.

Uji Heteroskedastisitas



Uji Autokorelasi

Model Summary ^b					
Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	Durbin-Watson
1	.726 ^a	.527	.513	1.10895	1.936

a. Predictors: (Constant), Penjualan, Modal Kerja

b. Dependent Variable: Laba Bersih

Uji Multikolinearitas

Coefficients ^a			
Model		Collinearity Statistics	
		Tolerance	VIF
1	Modal Kerja	.321	3.117
	Penjualan	.321	3.117

a. Dependent Variable: Laba Bersih

Regresi

Variables Entered/Removed ^a			
Model	Variables Entered	Variables Removed	Method
1	Penjualan, Modal Kerja ^b	.	Enter

a. Dependent Variable: Laba Bersih

b. All requested variables entered.

Model Summary ^b					
Model	R	R Square	Adjusted R Square	Std. Error of the Estimate	Durbin-Watson
1	.726 ^a	.527	.513	1.10895	1.936

a. Predictors: (Constant), Penjualan, Modal Kerja

b. Dependent Variable: Laba Bersih

ANOVA ^a						
Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	98.500	2	49.250	40.048	.000 ^b
	Residual	88.543	72	1.230		

Total	187.043	74			
-------	---------	----	--	--	--

a. Dependent Variable: Laba Bersih

b. Predictors: (Constant), Penjualan, Modal Kerja

		Coefficients ^a				
		Unstandardized Coefficients		Standardized Coefficients		
Model		B	Std. Error	Beta	t	Sig.
1	(Constant)	-1.938	.936		-2.071	.042
	Modal Kerja	.511	.388	.188	1.315	.193
	Penjualan	1.291	.328	.563	3.931	.000

a. Dependent Variable: Laba Bersih

Model 2 Setelah Dimoderasi Inflasi

Uji Normalitas

One-Sample Kolmogorov-Smirnov Test

		Unstandardize d Residual	
N		75	
Normal Parameters ^{a,b}	Mean	.0000000	
	Std. Deviation	1.08184723	
Most Extreme Differences	Absolute	.192	
	Positive	.150	
	Negative	-.192	
Test Statistic		.192	
Asymp. Sig. (2-tailed) ^c		.000	
Monte Carlo Sig. (2-tailed) ^d	Sig.	.000	
	99% Confidence Interval	Lower Bound	.000
		Upper Bound	.000

a. Test distribution is Normal.

b. Calculated from data.

c. Lilliefors Significance Correction.

d. Lilliefors' method based on 10000 Monte Carlo samples with starting seed 624387341.